

<https://helda.helsinki.fi>

---

## Draft genome sequences of three monokaryotic isolates of the white-rot basidiomycete fungus *Dichomitus squalens*

Casado López, Sara

2019-05-02

---

Casado López , S , Peng , M , Daly , P , Andreopoulos , B , Pangilinan , J , Lipzen , A , Riley , R , Ahrendt , S , Ng , V , Barry , K , Daum , C , Grigoriev , I V , Hilden , S K , Mäkelä , M R & de Vries , R P 2019 , ' Draft genome sequences of three monokaryotic isolates of the white-rot basidiomycete fungus *Dichomitus squalens* ' , Microbiology Resource Announcements , vol. 8 , no. 18 , e00264-19 . <https://doi.org/10.1128/MRA.00264-19>

---

<http://hdl.handle.net/10138/311844>

<https://doi.org/10.1128/MRA.00264-19>

---

acceptedVersion

---

*Downloaded from Helda, University of Helsinki institutional repository.*

*This is an electronic reprint of the original article.*

*This reprint may differ from the original in pagination and typographic detail.*

*Please cite the original version.*

**Draft genome sequences of three monokaryotic isolates of the white-rot basidiomycete  
fungus *Dichomitus squalens***

Sara Casado López<sup>a</sup>, Mao Peng<sup>a</sup>, Paul Daly<sup>a</sup>, Bill Andreopoulos<sup>b</sup>, Jasmyn Pangilinan<sup>b</sup>, Anna  
Lipzen<sup>b</sup>, Robert Riley<sup>b</sup>, Steven Ahrendt<sup>b</sup>, Vivian Ng<sup>b</sup>, Kerrie Barry<sup>b</sup>, Chris Daum<sup>b</sup>, Igor V.  
Grigoriev<sup>b</sup>, Kristiina S. Hildén<sup>c</sup>, Miia R. Mäkelä<sup>c#</sup>, Ronald P. de Vries<sup>a,c#</sup>,

<sup>a</sup>Fungal Physiology, Westerdijk Fungal Biodiversity Centre & Fungal Molecular Physiology,  
Utrecht University, Utrecht, The Netherlands

<sup>b</sup>US Department of Energy Joint Genome Institute, Walnut Creek, California 94598, USA

<sup>c</sup>Fungal Genetics and Biotechnology, Department of Microbiology, University of Helsinki,  
Helsinki, Finland

Running title: *Dichomitus squalens* genome sequences

#Address correspondence to Ronald de Vries, [r.devries@westerdijkinstitute.nl](mailto:r.devries@westerdijkinstitute.nl), Miia Mäkelä,  
[miia.r.makela@helsinki.fi](mailto:miia.r.makela@helsinki.fi)

21

## 22 **Abstract**

23 Here we report the draft genome sequences of three isolates of the wood white-rotting  
24 basidiomycete fungus *Dichomitus squalens*. These monokaryons were sequenced to provide  
25 more information on intraspecies genomic diversity of this fungus, also in relation to the  
26 previously sequenced genome of *D. squalens* LYAD-421 SS1.

27

## 28 **Main text**

29 *Dichomitus squalens* is a wood-decaying white-rot fungus commonly found in Europe, Asia and  
30 North America (1). It is mainly found on softwoods (2, 3), and has an extensive repertoire of  
31 lignocellulose-degrading enzymes (4-6). Two of the sequenced monokaryons, CBS463.89 and  
32 CBS 464.89, are derived from the well-studied Polish dikaryon FBCC312 (CBS432.34) (4, 6-  
33 11), while OM18370.1 is derived from the Finnish dikaryon OM18370 (CBS139088).

34 Strains were maintained on 2% (wt/vol) malt extract (ME) 1.5% (wt/vol) agar plates, from which  
35 four plugs (ø 5 mm) were used to inoculate stationary 50 mL 2% (wt/vol) ME liquid cultures,  
36 which were incubated at 28°C for five days. Genomic DNA was extracted from homogenized  
37 mycelium with extraction buffer (2% CTAB [N-cetyl-N,N,N-trimethylammonium

38 bromide], 100 mM Tris-HCl, 1.4 M NaCl, 20 mM EDTA, 0.2% β-mercaptoethanol) and purified  
39 with chloroform-isoamyl alcohol (24:1) (12). For RNA extraction, the isolates were pre-cultured  
40 on glycerol for 7 days (28°C) and transferred to solid-state cultures containing 2 g (dry weight)  
41 of Norway spruce wood sticks (2×0.2×0.2 cm) on top of 1% (wt/vol) water agar at 28°C for two

42 and four weeks (4). RNA extracts were layered over a 2 mL CsCl solution (5.7 M CsCl (Serva,  
43 Germany); 25 mM sodium citrate, pH 7.0; 0.5% *N*-lauroylsarcosine (Sigma, USA); 0.1 M  $\beta$ -  
44 mercaptoethanol (Sigma, USA)) in 13.2 mL polyallomer ultracentrifuge tubes (Beckman-  
45 Coulter, Brea, CA, USA) and centrifuged at 33,000 rpm for 21 h at 4°C in a Optima L-90 K  
46 ultracentrifuge, using the SW-41 Ti swinging bucket rotor (Beckman-Coulter, Brea, CA, USA).  
47 After centrifugation, supernatant was removed by, the tube was inverted and all but the bottom 1  
48 cm was sheared off. The RNA in the clear pellet was rinsed with 100  $\mu$ L of DEPC-treated water  
49 and then dissolved in 50  $\mu$ L of DEPC-treated water and stored at -80°C (13). The genomes were  
50 sequenced using Illumina platform and pairs of standard fragment (300 bp) and 4 Kbp long mate  
51 pair (LMP) libraries. Fragment libraries were produced from 100 ng gDNA sheared to 300 bp  
52 using the Covaris LE220 and size-selected using SPRI beads (Beckman Coulter). The fragments  
53 were treated with end-repair, A-tailing, and ligation of Illumina compatible adapters (IDT, Inc)  
54 using the Illumina library creation kit (KAPA biosystems). For LMP, 5  $\mu$ g of DNA was sheared  
55 using the Covaris g-TUBE™ (Covaris) and gel size selected for 4kb. The sheared DNA was  
56 treated with end repair and ligated with biotinylated loxP adapters and circularized by a Cre  
57 excision reaction (NEB). These were randomly sheared and treated as indicated for the fragment  
58 library, and enriched using eight PCR cycles for the final library.

59 For transcriptomes, which were used for genome annotations, stranded cDNA libraries were  
60 generated using the Illumina Truseq Stranded RNA LT kit. mRNA was purified using magnetic  
61 beads containing poly-T oligos, fragmented and reversed transcribed using random hexamers and  
62 SSII (Invitrogen) followed by second strand synthesis, and then treated with end-pair, A-tailing,  
63 adapter ligation, and eight PCR cycles.

The prepared libraries were quantified using KAPA Biosystem's next-generation sequencing library qPCR kit and run on a Roche LightCycler 480 real-time PCR instrument. The quantified libraries were then multiplexed with other libraries, and the pool of libraries was then prepared for sequencing on the Illumina HiSeq sequencing platform utilizing a TruSeq paired-end cluster kit, v4, and Illumina's cBot instrument to generate a clustered flow cell for sequencing. Sequencing of the flow cell was performed on the Illumina HiSeq2500 sequencer using HiSeq TruSeq SBS sequencing kits, v4, following a 2x150bp (2x100bp for LMP) indexed run recipe (14).

Illumina fastq files were QC filtered for artifact/process contamination. DNA reads were assembled AllPathsLG v. R49403 (15). For CBS463.89 lacking LMP, the initial assemblies of fragment data with Velvet v.1.2.07 (16) were used to create *in silico* long mate-pair libraries with insert 3000 +/- 300 bp. RNA reads were assembled using Rnnotator v.3.4.0 (17). All three genomes were annotated using the JGI Annotation pipeline v.1.9, which combines several ab initio, homology-based, and transcriptome-based gene predictors as well as tools and databases for functional annotation (18, 19).

All four genomes are highly similar in genome size and characteristics (Table 1). The improvement in sequencing methodology is reflected in the lower contig and gap number of the three new genomes compared to the older genome (LYAD-421 SS1). This data is highly useful to evaluate intraspecies genome variation in *D. squalens*.

#### **Data availability**

Genome assemblies and annotations are available via MycoCosm ([jgi.doe.gov/fungi](http://jgi.doe.gov/fungi); (18)). The data is deposited at DDBJ/EMBL/GenBank under BioProject/GenBank Accessions:

87 PRJNA334679/SELY000000000 (<https://www.ncbi.nlm.nih.gov/bioproject/PRJNA334679>),  
88 PRJNA334680/SELZ000000000 (<https://www.ncbi.nlm.nih.gov/bioproject/PRJNA334680/>),  
89 PRJNA334681/SELX000000000 (<https://www.ncbi.nlm.nih.gov/bioproject/PRJNA334681>).  
90

## 91 **Acknowledgements**

92 PD and MP were supported by a grant of the Netherlands Scientific Organization NWO  
93 824.15.023 to RPdV. The Academy of Finland grants no. 308284 and 297847 to MRM and KH,  
94 respectively, are acknowledged. We thank Ad Wiebenga for his assistance with the gDNA  
95 isolation. The work conducted by the U.S. Department of Energy Joint Genome Institute, a DOE  
96 Office of Science User Facility, was supported by the Office of Science of the U.S. Department  
97 of Energy under Contract No. DE-AC02-05CH11231. The authors declare no conflict of interest.

## 98 **References**

- 99 1. Andrews SR, Gill LS. 1943. Western red rot in immature Ponderosa pine in the  
100 Southwest. *J Forest* 41:565–573.
- 101 2. Blanchette RA, Otjen L, Carlson MC. 1987. Lignin distribution in cell walls of birch  
102 wood decayed by white rot basidiomycetes. *Phytopathol* 77:684-690.
- 103 3. Renvall P, Renvall T, Niemelä T. 1991. Basidiomycetes at the timberline in Lapland 2.  
104 An annotated checklist of the polypores of northeastern Finland. *Karstenia* 31:13-28.
- 105 4. Daly P, Casado López S, Peng M, Lancefield CS, Purvine SO, Kim YM, Zink EM,  
106 Dohnalkova A, Singan VR, Lipzen A, Dilworth D, Wang M, Ng V, Robinson E, Orr G,  
107 Baker SE, Bruijninx PCA, Hilden KS, Grigoriev IV, Mäkelä MR, de Vries RP. 2018.

*Dichomitus squalens* partially tailors its molecular responses to the composition of solid wood. Environ Microbiol 20:4141-4156.

5. Floudas D, Binder M, Riley R, Barry K, Blanchette RA, Henrissat B, Martinez AT, Otilar R, Spatafora JW, Yadav JS, Aerts A, Benoit I, Boyd A, Carlson A, Copeland A, Coutinho PM, de Vries RP, Ferreira P, Findley K, Foster B, Gaskell J, Glotzer D, Gorecki P, Heitman J, Hesse C, Hori C, Igarashi K, Jurgens JA, Kallen N, Kersten P, Kohler A, Kües U, Kumar TK, Kuo A, LaButti K, Larrondo LF, Lindquist E, Ling A, Lombard V, Lucas S, Lundell T, Martin R, McLaughlin DJ, Morgenstern I, Morin E, Murat C, Nagy LG, Nolan M, Ohm RA, Patyshakuliyeva A, et al. 2012. The Paleozoic origin of enzymatic lignin decomposition reconstructed from 31 fungal genomes. Science 336:1715-1719.

6. Rytioja J, Hilden K, Di Falco M, Zhou MM, Aguilar-Pontes MV, Sietiö O-M, Tsang A, de Vries RP, Mäkelä MR. 2017. The molecular response of the white-rot fungus *Dichomitus squalens* to wood and non-woody biomass as examined by transcriptome and exoproteome analyses. Environ Microbiol 19:1237-1250.

7. Casado López S, Peng M, Issak TY, Daly P, de Vries RP, Mäkelä MR. 2018. Induction of plant cell wall degrading CAZyme encoding genes by lignocellulose-derived monosaccharides and cellobiose in the white-rot fungus *Dichomitus squalens*. Appl Environ Microbiol 84:e00403-18.

8. Casado López S, Theelen B, Manserra S, Issak TY, Rytioja J, Mäkelä MR, de Vries RP. 2017. Functional diversity in *Dichomitus squalens* monokaryons. IMA Fungus 8:17-25.

- 129 9. Mäkelä MR, Sietiö O-M, de Vries RP, Timonen S, Hildén K. 2014. Oxalate-metabolising  
130 genes of the white-rot fungus *Dichomitus squalens* are differentially induced on wood  
131 and at high proton concentration. PLoS One 9:e87959.
- 132 10. Rytioja J, Hilden K, Hatakka A, Mäkelä MR. 2014. Transcriptional analysis of selected  
133 cellulose-acting enzymes encoding genes of the white-rot fungus *Dichomitus squalens* on  
134 spruce wood and microcrystalline cellulose. Fungal Genet Biol 72:91-98.
- 135 11. Rytioja J, Hilden K, Mäkinen S, Vehmaanperä J, Hatakka A, Mäkelä MR. 2015.  
136 Saccharification of lignocelluloses by carbohydrate active enzymes of the white rot  
137 fungus *Dichomitus squalens*. PLoS One 10:e0145166.
- 138 12. Hildén K, Martínez AT, Hatakka A, Lundell T. 2005. The two manganese peroxidases  
139 Pr-MnP2 and Pr-MnP3 of *Phlebia radiata*, a lignin-degrading basidiomycete, are  
140 phylogenetically and structurally divergent. Fungal Genet Biol 42:403-419.
- 141 13. Patyshakuliyeva A, Mäkelä MR, Sietiö O-M, de Vries RP, Hildén KS. 2014. An  
142 improved and reproducible protocol for the extraction of high quality fungal RNA from  
143 plant biomass substrates. Fungal Genet Biol 72:201-206.
- 144 14. Walker C, Ryu S, Na H, Zane M, LaButti K, Lipzen A, Haridas S, Barry K, Grigoriev IV,  
145 Quarterman J, Slininger P, Dien B, Trinh CT. 2018. Draft genome assemblies of five  
146 robust *Yarrowia lipolytica* strains exhibiting high lipid production, pentose sugar  
147 utilization, and sugar alcohol secretion from undetoxified lignocellulosic biomass  
148 hydrolysates. Microbiol Resour Announc 7:e01040-18.
- 149 15. Gnerre S, Maccallum I, Przybylski D, Ribeiro FJ, Burton JN, Walker BJ, Sharpe T, Hall  
150 G, Shea TP, Sykes S, Berlin AM, Aird D, Costello M, Daza R, Williams L, Nicol R,  
151 Gnirke A, Nusbaum C, Lander ES, Jaffe DB. 2011. High-quality draft assemblies of



mammalian genomes from massively parallel sequence data. Proc Natl Acad Sci U S A  
108:1513-1518.

16. Zerbino DR, Birney E. 2008. Velvet: algorithms for *de novo* short read assembly using de  
Bruijn graphs. Genome Res 18:821-9.

17. Martin J, Bruno VM, Fang Z, Meng X, Blow M, Zhang T, Sherlock G, Snyder M, Wang  
Z. 2010. Rnnotator: an automated de novo transcriptome assembly pipeline from stranded  
RNA-Seq reads. BMC Genomics 11:663.

18. Grigoriev IV, Nikitin R, Haridas S, Kuo A, Ohm R, Otilar R, Riley R, Salamov A, Zhao  
X, Korzeniewski F, Smirnova T, Nordberg H, Dubchak I, Shabalov I. 2014. MycoCosm  
portal: gearing up for 1000 fungal genomes. Nucleic Acids Res 42:D699-704.

19. Haridas S, Salamov A, Grigoriev IV. 2018. Fungal Genome Annotation. Methods Mol  
Biol. 1775:171-184.

167 **Table 1.** Genome statistics of the three *D. squalens* genomes CBS463.89, CBS464.89 and  
168 OM18370.1 compared to the previously sequenced genome of LYAD-421 SS1 (5).

	<b>CBS463.89</b>	<b>CBS464.89</b>	<b>OM18370.1</b>	<b>LYAD-421 SS1 (5)</b>
<b>Genome assembly size (Mbp)</b>	36.87	39.60	39.32	42.75
<b>Read coverage depth</b>	145x	118.7x	100.8x	50.63x
<b># of reads sequenced</b>	42.6M	32.7M	38.3M	7.7M
<b># of contigs</b>	1373	1147	1126	2852
<b># of scaffolds</b>	1259	467	439	542
<b>Scaffold N50</b>	134	44	39	16
<b>Scaffold L50 (Mbp)</b>	0.08	0.22	0.27	0.64
<b># of gaps</b>	114	680	687	1155
<b>% of scaffold length in gaps</b>	0.2%	2.5%	2.6%	7.7%
<b>Gene length (avg/median, bp)</b>	1,691/1,437	1,678/1,425	1,694/1,449	1,890/1,562
<b>Transcript length (avg/median, bp)</b>	1,370/1,140	1,358/1,128	1,365/1,150	1,484/1,213
<b>Exon length (avg/median, bp)</b>	259/158	259/158	256/157	254/152
<b>Intron length (avg/median, bp)</b>	76/61	77/60	78/61	86/61
<b>Protein length (avg/median, aa)</b>	387/314	382/311	388/319	419/345
<b># exons per gene (avg/median)</b>	5.3/4	5.25/4	5.34/4	5.84/4
<b># of gene models</b>	14,946	15,295	14,950	12,290
<b>GC content, %</b>	55.7	55.6	55.6	55.6